

SSITE SSITE

SPRING MIGRATION

North American Birds

A QUARTERLY JOURNAL OF ORNITHOLOGICAL RECORD PUBLISHED BY THE AMERICAN BIRDING ASSOCIATION

VOLUME 69: NO. 3, 2016 • MARCH THROUGH MAY 2015



THE NESTING SEASON

North American Birds

A QUARTERLY JOURNAL OF ORNITHOLOGICAL RECORD PUBLISHED BY THE AMERICAN BIRDING ASSOCIATION



VOLUME 69: NO. 4, 2016 • JUNE THROUGH JULY 2015

Northern California



Stephen C. Rottenborn Michael M. Rogers Jeff N. Davis Ed Pandolfino

SPRING

rought conditions continued to affect the entire Region in spring 2015, and temperatures were above average in March and early April before normalizing. Perhaps related to warm conditions, short-distance migrants such as Pacific-slope Flycatchers, Wilson's Warblers, and Hooded Orioles arrived earlier than average. At Southeast Farallon Island, the variety and number of migrant landbirds were exceptional, with several productive waves of migrants. The season's outstanding rarities included an apparent Red-backed Shrike x Turkestan Shrike hybrid, California's first Kelp Gull and second Tristram's Storm-Petrel, the Region's third and fourth Gull-billed Terns and fourth and fifth Wedge-rumped Storm-Petrels, and lingering Black Vulture, Rustic Bunting, and Brambling.

Abbreviations: C.B.R.C. (California B.R.C.); C.R.P. (Cosumnes River Preserve, *Sacramento*); C.V. (Central Valley); FI. (Southeast Farallon Island, *San Francisco*); H.R.S. (Hayward R.S., *Alameda*); P.R.N.S. (Point Reyes National Seashore, *Marin*); Point Blue (Point Blue Conservation Science); S.F. (San Francisco, not State Forest). Reports of exceptional vagrants submitted without documentation are not published. Documentation of C.B.R.C. review species will be forwarded to Tom Benson, Secretary, secretary@californiabirds.org.

WATERFOWL

Surprising numbers of geese lingered into

May. Greater White-fronted Geese were reported from seven counties through 31 May. An ad. blue-morph Snow Goose with 15 white morphs at the Sacramento N.W.R., Glenn 16 May was apparently the second-latest for the n. C.V. (ph. Steve Nord). A Ross's Goose lingering around Mountain View through 25 May (Eric Goodill, MMR, MJM) tied the late date for Santa Clara, and Cackling Geese were reported from four counties through 29 May. A Brant at the Salinas W.T.P., Monterey 14 May (KVV) was notable for its slightly inland location. Eurasian Wigeon in six counties through 31 May likewise represented more lingering birds than expected and included the latest for the n. C.V. at the Sacramento N.W.R., Glenn 18 May (†Bob MacMillan). Six Eurasian Teal were reported in five counties through 23 Apr. Male Tufted Ducks continued at Stafford Lake, Marin through 9 Mar (ph. Kevin Keirn, Donna Bray) and Lake Hennessey, Napa through 21 Mar (m.ob.); another was found at Abbotts Lagoon, Marin 24 Apr (Tate Snyder, AJ et al.). An unidentified eider was seen off Pigeon Point, San Mateo 13 May (RSTh). Of 3 Long-tailed Ducks, the latest was a lingering male at the Pajaro Dunes, Santa Cruz 6-9 May (NLv et al.). A female Hooded Merganser was on eggs in a nest box at Essex Pond, Humboldt 23 Apr (fide Elizabeth Feucht); another female along the Tuolumne River 16 May (HMR) raised thoughts of a possible first breeding record in Stanislaus. Inland Red-breasted Mergansers were found at Lake Camanche, Amador 13 Apr (DMr) and Crowley Lake, Mono 17 May (LKh).

LOONS THROUGH PELICANS

Red-throated and Pacific Loons are annual (albeit in low numbers) inland in early spring, but up to 2 Red-throated Loons at Lexington Reservoir 3-22 May (ph. Stephen Peterson et al.) provided Santa Clara's first May record, and a Pacific Loon at June Lake, Mono 23 May (ph. Phil Carnehl) was similarly late. Few Horned Grebes linger past Apr, so up to 3 at Alviso, Santa Clara 3-9 May (MMR, Gena Zolotar) and one at the Pajaro River, Santa Cruz 16 May (ph. Aaron Gabbe, Craig Fosdick) were notable. Santa Clara's wintering Red-necked Grebes at Calero Reservoir and Coyote Reservoir lingered through 9 Mar and 13 Apr, respectively (RPh, Andy Bradshaw, PLD, EFe et al.), but one at Lexington Reservoir 3-17 May (ph. Stephen Peterson et al.) was quite tardy. Dorris Reservoir hosted Modoc's second Red-necked Grebe 1-2 May (ph. SCR, JLx).

Laysan Albatrosses were well represented offshore, with a high of 9 seen from a repositioning cruise between *Monterey* and *Mendocino* 20 Mar (JCS, Tom Crabtree et al.). Even more exciting were the high numbers of Murphy's Petrels observed. Three off Sonoma 20 Mar (PEL, Nick Lethaby et al.) furnished a first county record. From late Apr through May, doubledigit counts were recorded on nine days, with highs of 68 between Monterey and Mendocino 27 Apr (Nathan Hentze), 63 from Mendocino to Del Norte 7 May (PEL, Gary Nunn, JCS, Monte Taylor et al.), and 54 in the vicinity of Davidson Seamount, Monterey 6-8 May (AdS et al.). The second record for F.I. was recorded 30 May (†Edward Jenkins). Twenty-two Hawaiian Petrels were recorded on these cruises 20 Mar-13 May, with a high of 9 off Mendocino and Humboldt 7 May (PEL, Gary Nunn, JCS, Monte Taylor et al.). Fewer Cook's Petrels were reported on repositioning cruises, with 17 seen on four dates 20 Mar-13 May, but 145 were seen on a research vessel 6-8 May around Davidson Seamount, Monterey (AdS et al.).

The difficulty of distinguishing Short-tailed Shearwaters from Sooties clouds our understanding of typical departure dates of the former, but it appears that only a few linger into Mar (and possibly early Apr). Two seen from shore at the entrance to Humboldt Bay, Humboldt 9 Mar (†RbF) were therefore noteworthy. The tail end of last winter's Black-vented Shearwater incursion provided the first spring record for S.F. in a decade, with 2 at Lands End 10 Mar (Collin Adams, Amanda Starbuck). Single Wedge-rumped Storm-Petrels mistnetted at F.I. 19 Apr and 20 May (ph. Point Blue) provided the fourth and fifth Regional records and the northernmost records to date; the previous three were from Monterey Bay. This species has typically been recorded off California only during warm-water conditions, and these records further exemplify the influx of southern seabirds that accompanied warmer water last winter. The significance of these records was eclipsed by California's second Tristram's Storm-Petrel, which was found dead (likely the victim of Burrowing Owl predation) at F.I. 18 Mar (ph. Point Blue); the first was also at F.I., on 22 Apr 2006.

A Blue-footed Booby at F.I. 22-24 Apr (Point Blue) provided a rare spring record, but one at the Klamath River mouth 20 Apr (†SMcA), furnishing Del Norte's second record, was far less expected so far north. Twenty-nine Brown Boobies more than doubled our previous spring total. At least 18 were at F.I., with high counts of 12 on 2 & 19 Mar (Point Blue); numbers there dwindled in Apr and early May but were bolstered by a few arrivals later in May. Elsewhere, 2 were in Monterey, 3 in San Mateo, 2 in S.F., 2 in Marin, and one in Humboldt. The Northern Gannet, now present for three years, was seen at F.I. most days (Point Blue), was detected around Alcatraz Island and the Golden Gate Channel, S.F. on a number of occasions 23 Apr+ (m.ob.), and provided *San Mateo's* first record at Daly City 25 May (PSar). A Brandt's Cormorant at Richmond 19 Apr (LKh, EPi, Jeff Hoppes) provided one of few recent spring records for *Contra Costa*, while a Pelagic Cormorant at Alviso, *Santa Clara* 30 May (MDo) may have been the same bird observed at this location sporadically over the past eight winters. The wintering American White Pelican wandered around coastal *Humboldt* and *Del Norte* until 21 Apr (m.ob.). A Brown Pelican was unusual (especially in spring) as far inland as Davis, *Yolo* 12 May (Susan Nishio).

BITTERNS THROUGH CRANES

Least Bitterns were reported in above-average numbers. Two were at Las Gallinas, Marin 16-18 May (Mark Dettling, Megan Elrod, Daphne Hatch), and up to 2 were at Campbell Cove, Sonoma 24 May+ (PBC, Mike Bumgarner, DN, m.ob.). Birds returned to Yolo Bypass W.A., Yolo 6 May+ (m.ob.), another was at the Cache Creek Conservancy, Yolo 16-20 May (Brent Campos et al.), and 4 were found at three locations in Sacramento 11 Mar-12 May (m.ob.). A Little Blue Heron courting a Cattle Egret at Natomas, Sacramento 8 May (Sarah Mayhew) was likely a returning bird but remained only a single day this year, unlike last summer when it spent 22 Jun-7 Jul at the same location. Up to 2 Cattle Egrets continued from winter through 16 Apr near Fort Dick, Del Norte (ADB).

Marin's Black Vulture continued in Bolinas through 10 May (RDiG, Alex Merritt, m.ob.). Another n. of Pacific Valley 14 May (†Greg Smith), providing the first record for Monterey, was deemed by the C.B.R.C. to be the same as a bird that has wandered the coast from Ventura to San Luis Obispo since 2009. A White-tailed Kite at Willow Creek W.A. 4 Mar (Peter Young, Jim Moore) was a rare find for Lassen. A Bald Eagle nest at Gallighan Slough held 2 young 6 Mar; both successfully fledged in early Jun, for the second successful nesting for this site and Santa Cruz. Hawkwatching from San Francisco on warm days with light to moderate offshore winds turned up 7 Broad-winged Hawks 18 Apr-3 May (HuC, DMo, m.ob.) and 2 Swainson's Hawks 30 Apr (HuC et al.). Both species were regarded as casual spring migrants prior to the recent regular monitoring of hilltop observation points. Farther south, northbound ad. Broad-wingeds were seen over Belmont, San Mateo 18 Apr (RSTh) and over Loma Prieta 19 Apr (second spring record for Santa Cruz and fifth for Santa Clara; ph. Pete Sole, DvWb et al.). Migrant Swainson's Hawks were seen over Belmont, San Mateo 25 & 27 Apr (both RSTh), over Quail Hollow Ranch, Santa Cruz 26 Apr (ph. AMR), and over Harkins Slough, Santa Cruz 15 May (ph. AMR, BRa, RRa). North of S.F., coastal migrants were found at Laguna de Santa Rosa, Sonoma 10 Apr (ScC) and e. of Willow Creek in Humboldt 3 May (TKz, †RbF, DCo, Casey Ryan). A pair nested again in Coyote Valley, Santa Clara, with nest building first observed 8 Apr (RPh, m.ob.).

Black Rails were well reported from Sacramento, with one at Stone Lakes N.W.R. 11-27 Mar (Mark Sawyer, m.ob.), one at Twitchell Island 17 Apr (Iryna Dronova), and up to 3 at C.R.P. 23 May-6 Jun (Breck Breckenridge, Julian Wood, JTr, m.ob.). Up to 4 were found at Alviso Marina C.P., Santa Clara 13 Mar-9 May (FV, m.ob.), with additional birds heard along nearby Alviso Slough 14 Apr+ (DvWb, WGB, MMR, MDo); another was heard farther n. at Coyote Hills R.P., Alameda 21 May (Jerry Ting). The Sandhill Crane at the Smith River bottoms, Del Norte was last seen 4 Apr (LzW, LPt). Lingering cranes in Humboldt included 2 near Ferndale 25 Apr (TMcK), 2 near McKinleyville 26 Apr (GSL, LPL), and 2 at Cock Robin Island 20 May (ph. Elizabeth Feucht et al.).

SHOREBIRDS

Continued drought resulted in 36-year-recordlow water levels at Lake Camanche, but this yielded near perfect conditions for shorebirds, particularly on the n. Amador side. In Amador, 5 Snowy Plovers 12 Apr-15 May (TEa, FS, Greg Hemig) may have been the first for the county, as was a Baird's Sandpiper 25 Apr (TEa, FS, ph. DMr, JCS, JLx et al.). Up to 3 Wilson's Phalaropes 7-13 May (TEa, ph. DMr, TS, Greg Hemig et al.) and up to 9 Marbled Godwits 21-25 Apr (ph. DMr, JLx et al.) provided the county's second and third records, respectively. Other notable shorebirds there included 40 Black-bellied Plovers, 4 American Avocets, 2 Black-necked Stilts, one Willet, 28 Whimbrels, 20 Long-billed Curlews, 10 Dunlin, and 3 Red-necked Phalaropes. Both Whimbrel and Long-billed Curlew were also seen in nearby Calaveras. Notable breeding records for American Avocets were obtained in Mendocino, where a pair incubated four eggs 30 Apr+ (SJS, CEV, RFK, m.ob.), and Santa Cruz, where three nests with eggs were found at College Lake 22 Apr-6 May (GKt et al.). Notable Black Oystercatchers inside S.F. Bay included 2 in Benicia, Solano 3 & 16 May (Ken Poerner, EPi), a nesting pair at Ballena Bay, Alameda (Vern Nelson, RJR, m.ob.), and Santa Clara's fourth, at Alviso 14 Apr (Jason St. Pierre) and 25 May+ (MMR, MJM, m.ob.). At least one American Golden-Plover was reported from Lake Earl and Lake Tolowa, Del Norte 21 May-4 Jun (ph. LBr). It was seen associating with a late Pacific Golden-Plover 25 May-4 Jun (ph. LBr).

The wintering Solitary Sandpipers reported last season continued through 9 Apr in Napa,

Napa (MBe, m.ob.) and through 8 Apr nw. of Millerton Lake, Madera (Nina Jones, m.ob.). Twenty-one migrant Solitaries reported from ten counties 16 Apr-9 May included Amador's second, at the Ione W.T.P. 25-26 Apr (TEa, FS, JCS, JLx, Greg Hemig), and a remarkable 5 at the Hayfork W.T.P. 30 Apr-4 May (fewer than five previous Trinity records, all of single birds; JLx, JSL). Humboldt's twelfth (and second in spring) Hudsonian Godwit graced Clam Beach 25-26 May (Mark Colwell, ph. SMcA, David Orluck, Deven Kammerichs-Berke, ph. RbF, m.ob.). Unusual shorebirds near Stratford, Kings included a Red Knot far inland 30 May and the only Stilt Sandpipers reported, up to 2 birds present 2-4 May (both MSy). Ruffs were at H.R.S. 18 Apr (RJR) and inland along Ostrom Rd., Yuba 28-30 Apr (JLx, JLa, ph. JCS, JSL, †BED, m.ob.). Single Sanderlings were far inland in Modoc, at Modoc N.W.R. 1 May and along Goose Lake Causeway 2-3 May (both SCR). Four Sanderlings at Alviso 25 May were the latest ever in spring for Santa Clara (ph. MMR, MJM). Besides the Amador bird noted above, Baird's Sandpipers were found near Kettleman City, Kings 9 Apr (2 birds; ph. JSy), along Abel Rd., Colusa 15 Apr (JCS), at Llano Seco North Central Valley W.M.A., Butte 19 Apr (MRg, Matt Forster), and at Colusa N.W.R., Colusa 24 Apr (JCS et al.). Spring Pectoral Sandpipers were found at Humboldt Bay N.W.R., Humboldt 20 Apr (Carrick Rice), Redding W.T.P., Shasta 24 Apr (Brooke McDonald), and s. of Corcoran, Kings 3 May (ph. MSy). Two of the 3 reported Semipalmated Sandpipers were the fifth and sixth in spring for San Mateo, at Ravenswood Trail, Menlo Park 23-26 Apr (RSTh et al., ph. MDeF, CHy) and Pilarcitos Creek mouth, Half Moon Bay 11 May (ph. MDeF, CHy); the third was at Colusa N.W.R., Colusa 24 Apr (†JCS).



This Gull-billed Tern, photographed on the last day of its 26 April–2 May 2015 stay south of Corcoran in Kings County, may have been a returning bird from last summer, which provided Northern California's second record. *Photograph by Mark Stacey.*

SKUAS THROUGH SKIMMERS

The only South Polar Skuas reported were off Eureka, Humboldt, with singles on 15 (RbF, SMcA, Mark Colwell et al.) & 21 May (PEL et al.). Long-tailed is the least frequently reported jaeger in spring; singles were off Monterey 6 May (PEL, Gary Nunn, JCS, Monte Taylor et al.), Trinidad Head, Humboldt 13 May (PEL et al.), and Eureka, Humboldt 15 May (RbF, SMcA, Mark Colwell et al.). Nine Scripps's Murrelets were seen on four dates, 7-21 May, between Mendocino and Del Norte. The total of 75 Sabine's Gulls, mostly alternate ads., between Monterey and Mendocino 20 Mar (JCS, Tom Crabtree et al.) was exceptional for such an early date. A good count of 5400 northbound Bonaparte's Gulls passed Pigeon Point, San Mateo 14 Apr (RSTh). Our only Laughing Gull was an ad. at Half Moon Bay, San Mateo 4-5 May (AJ et al.), while 8 Franklin's Gulls in six counties 29 Apr-25 May represented a modest showing. Among 3 Mew Gulls lingering into May, the latest was present through the period at Lake Merced, S.F. (m.ob.). Single Lesser Black-backed Gulls, both on 12 Mar, were at Lake Elizabeth, Alameda (ph. Jerry Ting) and Folsom Lake (third or fourth for Placer; RPz). Five Glaucous Gulls were recorded, the latest being at Half Moon Bay, San Mateo 11 May (MDeF, CHy). California's first Kelp Gull was well documented at Año Nuevo S.R., San Mateo 27 Apr (ph. Bob Siegel), Half Moon Bay, San Mateo 20 May (ph. AJ), and F.I. 24-29 May (ph. Point Blue); the same ad. was thought to be involved in all three appearances. This bird represented the first North American record w. of Colorado. A Gull-billed Tern s. of Corcoran, Kings 26 Apr-2 May (ph. MSy, m.ob.) may have been the same individual recorded there last May; one at the Pajaro River mouth 16 May (ph. JGa, Nicholas Kronick) was seen in both Santa Cruz and Monterey, providing a first record for each county. The Region had



The two Wedge-rumped Storm-Petrels mist-netted at Southeast Farallon Island in Marin County, California in spring 2015 (here 20 May) were the northernmost individuals recorded to date anywhere in the Pacific. *Photograph by Daniel Johnston.*

only two prior records. Six Black Terns at Lake Camanche 14-15 May (TEa, TS, Greg Hemig) were seen in both Calaveras and Amador, providing a first or second record for the latter county. The wintering Common Tern at Arcata Marsh, Humboldt lingered through 16 Mar (EE, RbF). Elegant Terns were early, with 2 at Bolinas Lagoon, Marin 20 Mar (ph. PP), 24 at Terrace Point, Santa Cruz 21 Apr (AMR), and 2 at Mountain View 20 Apr (Jason St. Pierre, Cole Jower), the latter providing the first Apr record for Santa Clara. Black Skimmers included the expected breeders in S.F. Bay and migrants in Monterey and Santa Cruz, whereas one at Ocean Beach 10 May (HuC, PSar) provided only the fifth S.F. record.

DOVES THROUGH THRUSHES

The Band-tailed Pigeon incursion in the lowlands continued well into spring, highlighted by what may have been Modesto's first record, involving 2 birds at Tuolumne River R.P., *Stanislaus* 19 Apr (HMR). The last lowland report of the season was of a single bird at Oroville, *Butte* 27 May (Carol Anderson). Our only White-winged Dove was at San Mateo, *San Mateo* 28 May (ph. Mike Aicardi). A Great Gray Owl with a broken wing found at Lake Mary near Mammoth Lakes 22 May (*fide* KNN) provided only the second or third record for *Mono*.

Astounding numbers of Anna's Hummingbirds visited multiple feeders at a residence in Carmel Valley, *Monterey* during the period; the peak estimate of more than **900** on 30 Apr (PFw), based on timed counts and presumed turnover rates from the literature, recalled densities at certain feeders in the mountains of Arizona. This total eclipsed our previous highest one-day estimate of 150 at Springville, *Tulare* 6 Sep 1980. A female Costa's Hummingbird at Napa 10 Apr (Craig Cummings) established about the fifth record of that species for *Napa*. A total of 38 ad. male Rufous Hummingbirds

> at Summit Ridge 11 Apr (JGa, AMR, Nicholas Kronick) represented a strong showing for Santa Cruz. Also impressive were 20 ad. males at nearby Loma Prieta Ave. on the Santa Cruz/Santa Clara border 17 Apr (JGa, Kevin Condon, Nicholas Kronick). A high count of 7 Calliope Hummingbirds at Summit Ridge 17 Apr (JGa, Nicholas Knonick, m.ob.) more than doubled the previous record of 3 for any spring season in Santa Cruz. Likewise, 10 in Monterey during the period established a new seasonal high count for that county (fide DR), and 5 at Dinosaur Point Rd. 1 May (MMR, RGJ, RFu) set a new single-locality high count for Merced.

The fall-winter invasion of Lewis's and Acorn Woodpeckers fizzled out in spring, with no especially notable numbers or occurrences. A Williamson's Sapsucker along Hwy. 36 on the Shasta/Trinity border 4 May (JSL) provided the first spring record for Trinity. Although this species may be resident in the ne. portion of the county, this bird was about 100 km sw. of that area. Six Yellow-bellied Sapsuckers included 5 coastal (one in Marin, 2 in San Mateo, one in Santa Cruz, and one in Monterey) and one inland at O'Neill Forebay, Merced 12 Mar (PDu, FrO), continuing from winter. Similarly, 8 Red-naped Sapsuckers involved 5 coastal and bayside birds (one in Solano, one in Alameda, 2 in Santa Clara, and one in Santa Cruz) and 3 inland (2 in Lassen and one in Merced). An unusual movement of Pileated Woodpeckers brought one to the Presidio 16 Apr (the third or fourth for S.F.; HuC, BFi, LKh), 3 to Contra Costa, and 2 to Monterey. All but one occurred within the period of 14-29 Apr; the outlier was one at Lake Anza, Contra Costa 17 May (Christine Woo).

Large numbers of Hammond's Flycatchers moved through Santa Clara, with 10 reported 3 Apr-23 May (twice the average fide WGB), and Mono, with 39 reported 26 Apr-31 May (a new seasonal high count fide KNN). All but one of Napa's records of Gray Flycatcher have come from Wild Horse Ridge; one bird there 27 Apr and another there 1 May (ph. MBe) provided the fifth and sixth records for the county. Black Phoebes continue to increase their presence in Lassen, with multiple reports during the season, including from as far e. as Biscar Reservoir and Honey Lake (fide KAb). Typical for the season, 2 Eastern Phoebes were reported: one at Pescadero, San Mateo 8 Mar (LFl), and another at Bolinas Lagoon, Marin 18 Mar (JsC). Sonoma's first Vermilion Flycatcher was a female along the Santa Rosa Creek trail in Santa Rosa 20 Apr (DgSh). Another at Oasis Ranch 22-23 May (PJM, ADeM) established the third spring record for Mono. Tropical Kingbirds, most continuing from winter, included one in S.F., 2 in San Mateo, one in Contra Costa, and 2 in Santa Cruz. Scarce anywhere along the Region's immediate coast, a Cassin's Kingbird stopped in at the lighthouse at Point Reyes N.S., Marin 30 Apr (ph. DSg). Our only Eastern Kingbird visited F.I. 25-27 May (Point Blue).

The most thought-provoking find of the season was a "brown" shrike in Manchester, *Mendocino* 5 Mar–22 Apr (ph. Alison Cebula, ph. m.ob.). Not clearly a Brown Shrike (*Lanius cristatus*), for which there are three previous records for California, this bird most closely resembled a **Red-backed Shrike** (*Lanius collurio*) x Turkestan Shrike (*L. phoenicuroides*) hybrid (see North American Birds 69: 4-35).

Not only are there no previous records of this hybrid for North America, there are no records of these Old World parent species either. Two Least Bell's Vireos were reported: one singing w. of Corcoran, Kings 9-31 May (ph. MSy, ph. m.ob.) and another at Belmont, San Mateo 14 May (RSTh). A total of 25 Warbling Vireos at Pacheco S.P. 9 May (MMR, PDu) represented the largest pulse of migrants of that species ever reported from one location in Merced. Red-eyed Vireo is less than annual in spring, so the total of 4 this spring was remarkable. Singles were at Golden Gate Park, S.F. 9 May (ASH), Putah Creek, Yolo 18 May (AEn), Point Reyes N.S., Marin 28 May (DSg), and Tunitas Creek Rd., San Mateo 29 May (RSTh).

The invasion of Steller's Jays on the C.V. floor that began last fall continued into May in the n. C.V. with numerous reports from Sutter, Yuba, Butte, Glenn, and Tehama (fide BED). Goldencrowned Kinglets nested at the Strybing Arboretum in Golden Gate Park during the period (m.ob.), apparently for the second consecutive year, representing the first confirmed breeding records for S.F. Six Townsend's Solitaires were found outside areas of residency during the period, with one in Tehama, 2 in Napa, one in S.F., one in San Mateo, and one in Contra Costa. Small numbers of residual Varied Thrush from the massive fall-winter invasion lingered in the C.V. and along the coast through spring, with some in Monterey even hanging on into Jun (fide DR).

THRASHERS THROUGH WARBLERS

Our only Gray Catbird report was of a briefly glimpsed bird during a late snowstorm near McGee Creek, Mono 15 May (ADeM). Six outof-range Sage Thrashers included 4 along the coast, one in the C.V., and one at n. Lake Tahoe, Placer. Lapland Longspurs were at Ferndale bottoms, Humboldt 1 Mar (AxL), on F.I. 12 May (Point Blue), and at Lake Tolowa, Del Norte 22 May (LBr). Two Chestnut-collared Longspurs continued from winter on Bear River Ridge, Humboldt through 4 Apr (TKz, m.ob.). An Ovenbird at Charleston Road marsh, Santa Clara was a winter bird that barely continued into spring 1 Mar (AV). Ovenbirds were also in Cottonwood Canyon, Mono 23-26 May (PJM, m.ob.) and on F.I. 24 May+ (Point Blue). A total of 6 Northern Waterthrushes was more than double our average and included a singing bird likely returning for its third spring near Lake Almanor, Plumas 9 May (RDB, CPD). Bluewinged Warblers are reported in our Region at a rate of about one every five years, so the male that landed on a boat 3 km off Moss Landing, Monterey 24 May (Kate Spencer, ph. Don Bacchus) was a very good find. All but one of our 9 Black-and-white Warblers (twice our average) were in coastal or Bay Area counties. The exception was along McGee Creek, Mono 18 May (CMc). A Prothonotary Warbler at Bodega Bay, Sonoma 30 May (Mike Heffernon) and 3 Tennessee Warblers at Point Reyes, Marin 29-31 May (Everett Clark, LKh, m.ob.) all arrived just in time to provide late spring records. The Lucy's Warbler at Bodega Bay, Sonoma 9-14 Mar (DeH, RAR, m.ob.) provided our first spring record since 2009. A Kentucky Warbler at Point Reyes, Marin 28 May sang a bit and even posed briefly for a photograph (ph. DSg). Our only Hooded Warbler was on F.I. 14-16 May (Point Blue). A male American Redstart in Philo, Mendocino 1 Mar-19Apr (TDB, m.ob.) sang throughout his visit. Our other redstart was a female in a Pacific Grove, Monterey backyard 25-29 May (ph. RC, ph. DR, m.ob.). We average more than seven Northern Parula reports in spring and have received no fewer than two for at least the past 25 springs. Therefore, our lone parula at Point Reyes, Marin 30 May (Mary Ann & Mark Butler, Roger Harshaw) marked a season low for the Region. Magnolia Warblers were in Cottonwood Canyon, Mono 24 May (Meg Andrews) and at Pacific Grove, Monterey 27 May (ph. RC, ph. MiR). Three Chestnut-sided Warblers provided our best spring total for this species in more than 25 years, with males at Mendocino Highlands S.P., Mendocino 25 May (M. Gertz), at Willow Glen, Santa Clara 26 May (Brooke Miller; first spring Santa Clara record), and on F.I. 31 May (Point Blue). We tallied 16 Palm Warblers, twice our average, and this included one Yellow Palm Warbler on F.I. 25 Apr (Point Blue). A singing Yellow-throated Warbler in Bodega Bay 27 May (v.r. DN) furnished a first for Sonoma.

SPARROWS THROUGH FINCHES

Two of the season's 3 Clay-colored Sparrows were in Mono, one at South Tufa 1 May (†George Chisman) and another at Bridgeport 22 May (KNN, JLD), with a third bird on F.I. 18 Apr (Point Blue). Black-chinned Sparrows are sparsely distributed and irregular breeders in the Region. This spring they continued for the fourth year near Loma Prieta in both Santa Clara and Santa Cruz, with singing birds arriving by 4 Apr (JGa, AMR, m.ob.). In contrast, they have apparently not bred on Mount Diablo, Contra Costa for several years, and this year's only Black-chinned Sparrow reported there 19 Apr did not remain (LKh, EPi, Jeff Hoppes). Eleven singing Black-chinned Sparrows were along Bear Mountain Rd. in Henry Coe S.P., Santa Clara 16 May (MMR). Two Black-throated Sparrows were at n. Lake Tahoe, Placer 30 Apr-2 May (ScD). A Nelson's Sparrow continued from winter through 20 Mar at Arrowhead Marsh, Alameda (Mark Rauzon). The season's

latest lingering White-throated Sparrow was at Marina, *Monterey* until 22 May (SRv). Also continuing from winter were 4 Harris's Sparrows, one each at Red Bluff, *Tehama* 18 Mar (MSw), Salmon Creek, *Sonoma* 10 Apr (LHg), Modesto, *Stanislaus* 21 Apr (Tom & Anne Myers), and Arcata, *Humboldt* 30 Apr (GAB). Our winter Rustic Bunting in *S.F.* continued through 31 Mar (m.ob.).

Eleven Summer Tanagers included a remarkable 3 at Oasis Ranch, Mono 22-23 May (PJM, ADeM, CMc). Twenty-two Rose-breasted Grosbeaks provided a good showing, as we average fewer than 14. A Blue Grosbeak was at Spenceville W.A. 10 May (ECB, RCD, John Lace), the same location that hosted Nevada's first last year. Indigo Buntings were present in typical numbers (8 vs. an average of 7), but there was nothing average about an ad. male Painted Bunting at Bridgeport, Mono 20 May (ph. Tim Taylor, Ryan Spaulding). Tricolored Blackbirds are in significant long-term decline in the state, so it was encouraging to receive reports of new or re-colonized breeding locations. Up to 30 Tricoloreds along the Carson River 3-21 May (TEa, LKh, m.ob.) provided Alpine's first documented record, a new colony was found 28 Mar near Fall River, Shasta (B&CY, KAb, DMr), and nesting was confirmed at a site in Potter Valley, Mendocino 8 May (GEC), where the birds have been absent for about 10 years. Orchard Orioles were at the Presdio, S.F. 16 Apr (HuC, LKh, BFi), on F.I. 17-18 May (Point Blue), and at Belmont, San Mateo 31 May+ (RSTh, LBt). An unexpected Hooded Oriole at s. Lake Tahoe, El Dorado 7-10 Apr (Bob Sweatt, MMy) provided a first for the Lake Tahoe Basin (El Dorado, Placer, and Nevada). Baltimore Orioles included a bird continuing from winter at San Mateo, San Mateo through 6 Mar (Merry Haverman), a male at Capitola, Santa Cruz 8 Apr (Judy Donaldson), and a male and female in different locations in Bodega Bay, Sonoma 26-29 May (RAR, DN, m.ob.). An oriole at Westgate Landing C.P., San Joaquin 15-31 May (Pat & Dave Croft, DGY, JLD, m.ob.) showed clear signs of being a Baltimore x Bullock's hybrid. A Scott's Oriole was at Paradise, Mono 4-18 Apr (DHo). The Brambling found in winter in Arcata, Humboldt continued through 12 Mar (Cindy Kuttner). A Cassin's Finch visited a feeder in S.F. 20 Apr (PSar).

Cited observers (county subregional editors in boldface): Ken Able (Lassen), Alan D. Barron, Leonie Batkin, Murray Berner (Napa), William G. Bousman (Santa Clara), Tim D. Bray, Lucas Brug, Ryan D. Burnett, Scott Carey, Rita Carratello, George E. Chaniot, Josiah Clark, Peter B. Colasanti, Daryl Coldren (Humboldt and Del Norte), Chris Conard (Sacramento), Hugh Cotter, Rudy C. Darling (Nevada), Jeff N. Davis (Madera), Malia DeFelice, Al DeMartini, Bruce E. Deuel (n. C.V. counties), Scott Dietrich, Ryan DiGaudio, Colin P. Dillingham (Plumas and Sierra), Matthew Dodder, Peggy L. Don, Jon L. Dunn, Pete Dunten, Todd Easterla, Elias Elias, Andrew Engilis, Paul Fenwick, Eric Feuss, Brian Fitch, Leslie Flint, Rick Fournier (RF), Rob Fowler (RbF; Humboldt and Del Norte), Rob Furrow (RFu), John Garrett, Steve Gerow (Santa Cruz), John Harris (Stanislaus), Karen Havlena (Mendocino), Chris Hayward, Alan S. Hopkins, Debbie House, Lisa Hug, John E. Hunter (Trinity), Alvaro Jaramillo, Richard G. Jeffers, Logan Kahle (Contra Costa), Robert J. Keiffer (Mendocino), Gary Kittleson, Tony Kurz, Alexandra Lamb, Jim Laughlin, Paul E. Lehman, Robin L. C. Leong (Solano), Gary S. Lester, Lauren P. Lester, Nick Levendosky, Jim Lomax, John S. Luther, Michael J. Mammoser, Don Marsh, Dan J. Maxwell (Marin), Sean McAllister, Chris Mc-Creedy, Tristan McKee, Peter J. Metropulos (San Mateo), Dominik Mosur (San Francisco), Martin Meyers, Dan Nelson, Kristie N. Nelson (Mono), Frances Oliver (San Joaquin), Ed Pandolfino (Placer), Ryan Phillips, Eric Pilotte, Linda Pittman, Gary W. Potter (Fresno), Ron Pozzi, Peter Pyle, Bernadette Ramer, Robert Ramer, Harold M. Reeve, Robert J. Richmond (Alameda), Michael Rieser, Alex M. Rinkert, Don Roberson (Monterey), Michael M. Rogers (MMR), Michael Rogner (MRg), Stephen C. Rottenborn, Steve Rovell, Ruth A. Rudesill (Sonoma), Paul Saraceni, Adam Searcy, Jeff Seay (Kings), Doug Shaw, Daniel Singer, Mark Stacy, John C. Sterling (Modoc, Alpine, Calaveras, and Yolo), Fritz Steurer, Tim Steurer (El Dorado and Amador), Steve J. Stump, Steven D. Summers (Tulare), Michele Swartout, Ronald S. Thorn, Jim Tietz (F.I.), John Trochet, Steve Umland (Tuolumne), Frank Vanslager, Chuck E. Vaughn, Ann Verdi, Kent Van Vuren (Merced), Dave Weber, Liz West, Jerry R. White (Lake), David G. Yee, Bob & Carol Yutzy (Shasta). Many more observers are not specifically cited; all are appreciated.

SUMMER

earshore sea surface temperatures were above average in June and near record-high values in July. At Southeast Farallon Island, breeding success by most seabird species was about average, despite reduced numbers and a delayed start to breeding. Warm-water conditions are usually associated with reduced seabird breeding success, but strong upwelling in early spring apparently benefitted the local food web, which also brought Sooty Shearwaters, Elegant Terns, and Sabine's Gulls to nearshore areas in unusually large numbers. Notable rarities this season included the continuing Black Vulture, a Curlew Sandpiper, and a Little Stint. Significant breeding records included the southernmost breeding by Harlequin Ducks in North America.

WATERFOWL THROUGH PELICANS

Brant inside S.F. Bay included 4 at Brooks Island, Contra Costa 27 Jul (LKh, CDu), where they have been fairly regular in recent years, as well as single birds at Aquatic Park, S.F. 10 Jul (ph. Carlo Arreglo) and Coyote Point, San Mateo 24-28 Jul (RSTh, AME). Two green-collared Trumpeter Swans from the introduced Oregon population found s. of Alturas, Modoc 15 Jul (SCR) provided only the second summer record for the Region; might pairs from that population eventually breed in California? A very late Eurasian Wigeon lingered from spring at Arcata Marsh, Humboldt through 3 Jun (m.ob.). Two female Ring-necked Ducks with about 14 ducklings near Truckee 7 Jul (ph. WGB) were thought to provide the first breeding record for Nevada. A single Hooded Merganser duckling with them was likely a result of egg dumping. Lesser Scaup bred again at H.R.S., with the first downy young noted 5 Jul (RJR), but no breeding evidence was noted in Santa Clara for the second year in a row. A female Harlequin Duck with 5 ducklings along the San Joaquin River near Balloon Dome 11 Jul (Ian Allison, ph. Kearey Allison) provided only the second breeding record for Madera, and the first in 91 years! This likely constitutes the southernmost breeding location for North America and presumably the world. Unseasonable scoters inside S.F. Bay (all in San Mateo) included a Whitewinged at Redwood City 21 Jun (female; RSTh), another at Coyote Point 29 Jun (male; ph. Gena Zolotar), and a Black at the latter location 5 Jul (male; ph. Warren & Marilyn Regelmann). Unusual far inland, and especially so in summer, an ad. male Red-breasted Merganser stayed at Bridgeport Reservoir, Mono 19 Jun-6 Jul (PJM et al., ph. RCP).

A Red-throated Loon at Lexington Reservoir lingered from spring through 2 Jun (MMR), furnishing *Santa Clara* with its first summer record. A concentration of 150 Pied-billed Grebes at Clifton Court Forebay 27 Jul (LKh, CDu) provided one of the highest counts ever for *Contra Costa*. Rare summer Horned Grebes were at Crowley Lake, *Mono* 15 Jun (CMc), Mountain View, *Santa Clara* 16 Jun (Jasen Liu), and Eagle Lake, *Lassen* 28 Jul (DPA). Eared Grebes nested again at H.R.S., where they have bred each year since 1999 (RJR); this is the most regular S.F. Bay-area breeding location. In contrast, bayside nesting by Clark's Grebes has been documented only since 2011. This season, at least three nests were present in a former Mountain View, Santa Clara salt pond 6 Jun+ (MMR, MJM, RGJ), including one built on an old hunting blind. A cruise on 12 Jul produced a Hawaiian Petrel off Humboldt and 8 Ashy Storm-Petrels (3 off Del Norte and 5 off Mendocino; Jeff Gilligan, Owen Schmidt). A Manx Shearwater was seen from Moss Beach, San Mateo 31 Jul (RSTh); we average 1-2 each summer. The sulid trio of Northern Gannet, Blue-footed Booby, and Brown Booby was once again recorded at F.I.; a Bluefooted was seen 19-30 Jul, up to 5 Brown Boobies were present throughout the period (with the high of 5 on 4 Jun), and the gannet was observed sporadically both at FI. and at Alcatraz Island, S.F. (Point Blue, m.ob.). Otherwise, the only booby was a Brown at Half Moon Bay, San Mateo 26 Jun (MDeF). A Brandt's Cormorant at Miller/Knox R.S. 11 Jul (LKh, CDu) provided Contra Costa's first summer record since 2008. American White Pelicans are scarce in most coastal areas; concentrations of 90+ at Bodega Harbor, Sonoma 3 Jun (Len Blumin), 280 at Struve Slough, Santa Cruz 18 Jul (DSy), and up to 200 at Elkhorn Slough, Monterey throughout the period (m.ob.) were unusually large. One was at Roberts Lake, Monterey 22 Jul (ph. SLn, ph. RC, DR); there are fewer than 10 Monterey Peninsula records. A Brown Pelican at West Sacramento 12 Jun (Aaron Haiman) provided one of few Yolo records.

BITTERNS THROUGH CRANES

Least Bitterns continued to be reported in above-average numbers. Up to 2 continued at Las Gallinas, *Marin* through the period (ph. m.ob.), one continued at Campbell Cove, *Sonoma* through 30 Jun (DN, JLx, m.ob.), up to 3 continued at Yolo Bypass W.A., *Yolo* 29 Jun–19 Jul (ph. m.ob.), and one was found again at Mather Lake, *Sacramento* 12 Jun (Phil Lutz). In addition, up to 3 birds were found at Clear Lake 13 Jun–24 Jul (Greg Guisti, Brad Barnwell), the first found in *Lake* in two years.

Marin's Black Vulture was seen again over Chimney Rock at P.R.N.S. 18 Jul (ph. Carlo Arreglo). Bald Eagles nested at Crystal Springs Reservoir, San Mateo for the third year in a row, successfully raising 2 young this year. An unseasonable Broad-winged Hawk heading n. over P.R.N.S. 4 Jun (ph. DSg) apparently provided the first summer record for the Region. Swainson's Hawks nested successfully again near the Napa County Airport, Napa (one fledgling at nest 31 Jul; MBe) and along Coyote Creek at Coyote, Santa Clara (one nearly fledged nestling 16-24 Jul; WGB, RPh).

Up to 4 Yellow Rails were found again s. of Eagle Lake, *Lassen* 7 Jun–4 Jul (m.ob.). Black Rails continued to be found along Alviso Slough and at Alviso Marina C.P., *Santa Clara*, with a high count of 4 detected 4 Jul (SCR). Common Gallinules bred successfully again in *Mendocino*, with at least 2 young detected at Ukiah 20 Jun (SJS). Young from the first county breeding attempt at the same location in 2012 were thought to have perished when the pond was cleared of all wetland vegetation. A Sandhill Crane was seen again at the Smith River bottoms 2 Jul (Maureen Kevany-Jahn), and another was heard at Tolowa Dunes S.P. 15 Jul (both *Del Norte*; TKz).

SHOREBIRDS

The nesting American Avocets at the Ukiah W.T.P., *Mendocino* hatched 3 young 3 Jun and remained through 16 Jul (ph. SJS, m.ob.). Notable S.F. Bay Black Oystercatchers continued from spring with 2 at Benicia, *Solano* 7 Jul (FrO, LPt), up to 6 (including 2 young) at Ballena Bay, *Alameda* (RJR, m.ob.), and *Santa Clara*'s fourth at Alviso through 22 Jun (Bill Pelletier). An alternate-plumaged American Golden-Plover was found at Arcata Bay, *Humboldt* 27 Jun–14 Jul (ph., †TMcK, EE, ph. RbF, m.ob.). Two other alternate-plumaged golden-plovers there 6 Jul (EE, TMcK) were not identified to species.

Wandering Long-billed Curlews ended up in unusual locations. One at Clear Lake 3 Jul (ph. FHa) was only the sixth for Lake, up to 2 at Boca Reservoir 12-13 Jul (ph. SRs, DRs, m.ob.) provided a rare record for Nevada, and singles made it to F.I. 3 & 14-15 Jul (Point Blue). A Ruddy Turnstone at Clear Lake S.P. 30 Jul (JRW) provided only the second record for Lake, following the first at the same location 14 years earlier. Another far inland near Corcoran, Kings surprisingly lingered 13 Jun–18 Jul (ph. MSy). Reports of white-necked molting ad. male Ruffs retaining long white chest plumes at the Davis Wetlands, Yolo 29 Jun (†LJE, †John Robinson) and near Corcoran, Kings 11 Jul+ (ph. MSy) probably involved the same bird. Santa Clara's fourth Curlew Sandpiper was a basic-plumaged bird at Alviso 5 Jul (†MJM). The Region's fifteenth Little Stint, and second for Kings, remained near Corcoran 11-19 Jul (ph. MSy). Among 26+ Semipalmated Sandpipers 30 Jun+ was the third or fourth for Napa, at American Canyon Wetlands 24 Jul (MBe, ph. Ken Skolnik, m.ob.). Early returning Wilson's Snipe included a bird at Zmudowski S.B., Monterey 19 Jul (ph. Michael Rieser), up to 2 birds at Mountain View 28 Jul+ (earliest fall arrival for Santa Clara by three days; ph. Bob Bolles et al.), one at the Yolo Bypass W.A., Yolo 28 Jul (ph., fide eBird), one at the Ukiah W.T.P., Mendocino 30 Jul (DAB, ph. SJS), and one at the Madera W.T.P., Madera 31 Jul (ph. GaW, Rick Saxton). Also notable away from known breeding areas were two courting pairs of snipe along Bentley Rd., *Stanislaus* 22 Jun (HMR) and a single bird along Dairy Rd., *Yuba* 27 Jun (ph. Marge & Don Thornton).

JAEGERS THROUGH TERNS

Good counts of 22 Pomarine and 30 Parasitic Jaegers harassed gulls and terns off Pajaro Dunes, Santa Cruz 27 Jul (AMR, NLv). Sabine's Gulls (about 95% ads.) staged an unprecedented nearshore incursion in Monterey Bay 18 Jul+, with high counts of 1575 at Pajaro Dunes, Santa Cruz 27 Jul (AMR, NLv) and 1870 in a 1.75-hour count from Point Pinos, Monterey 28 Jul (BLS, DR, RC). Although water temperatures were high (typically associated with low food supplies) throughout Monterey Bay in Jul, prey species such as Northern Anchovy were concentrated inshore, resulting in these record congregations. Rarer gulls included a first-cycle Little near Stratford, Kings 22 Jun-12 Jul (ph. MSy), a second-cycle Laughing at Bodega Harbor, Sonoma 22-23 Jul (Kirsten Lindquist), and a first-cycle Glaucous at Crescent City Harbor, Del Norte 23 Jun (ADB). Eight Franklin's Gulls scattered among five counties represented a typical summer showing for recent years. Mew Gulls are rare in summer, so singles at Alviso, Santa Clara 13 Jul (WGB, MJM) and at Lake Merced, S.F. throughout the period (m.ob.) were noteworthy. Although Western Gull records in the C.V. have increased in recent years, most such occurrences have been in winter, so a firstsummer bird at El Rico, Kings 7 Jun (ph. MSy) was unexpected. The number of breeding Western Gulls at F.I. was 30% lower than last year, reaching the lowest breeding population in 45 years of monitoring (Point Blue). Chick survival was low early due to starvation and intraspecific predation but improved in Jul. California Gulls once again attempted breeding at F.I., but none were successful, primarily owing to nest destruction by loafing sea lions (Point Blue).

Foraging Least Terns from colonies at Huichica Creek, Napa and the Montezuma Wetlands, Solano were observed in a number of bayside areas in those two counties. Ninety at Huichica Creek colonies 15 Jul (MBe) provided a high count for that location. Least Terns nested again at the Sacramento W.T.P., Sacramento, with a pair first detected 5 Jul and 2 chicks hatching at the end of the month (CCo, KLL, Steve Scott et al.). This species nested here 2008-2012, made a brief appearance in 2013, and was absent last year. Extralimital Least Terns were reported more widely than ever before. One at the Chico W.T.P., Butte 10 Jun-16 Jul (MRg, JLw, ph. Matt Forster) was likely one of last summer's birds returning. A Least Tern at Crowley Lake 14 Jun (DHo, Chris Allen) provided Mono with its third record, whereas singles at the Madera

W.T.P. (George Folsom) and Woodland W.T.P. (ph. Keith Bailey), both on 17 Jul, provided fourth records for Madera and Yolo, respectively. Individuals at Pajaro Dunes, Santa Cruz 5 Jul (Aaron Gabbe) and 28 Jul (Jonathan Vargas) may have been southbound migrants from S.F. Bay colonies. Seventeen Black Terns on 19 Jul provided a good count at Thermalito Afterbay, Butte (FrO, ClH, JRow), where the species occurs as a regular migrant, whereas singles at H.R.S. 20 Jun (RJR), Sunnyvale W.T.P., Santa Clara 20 Jul (Vivek & Leena Khanzode), and Franks Tract, Contra Costa 28 Jul (†ES) were in areas of less frequent occurrence. Single ad. Arctic Terns at H.R.S. 7 Jun (RJR) and Redwood Shores, San Mateo 6 Jul (RSTh) were rare inside S.F. Bay, particularly in summer, and may have represented the same individual. More than 30 pairs of Forster's Terns nested at Woodward Reservoir, Stanislaus this summer (JHH et al.); this species is a scarce breeder in the San Joaquin Valley and had not nested at Woodward Reservoir since 1999. Elegant Terns invaded coastal areas in large numbers for the second consecutive summer, beginning in late Jun; by late Jul, thousands were present, especially in Monterey Bay, with large numbers (including many juvs.) foraging on anchovies in inshore waters (DR, m.ob.).

DOVES THROUGH THRUSHES

Two Band-tailed Pigeons at Oroville, Butte 30 Jun (Carol Anderson) were unexpected at that location and presumably lingered from the fall-spring incursion. Providing the first record for the greater Sacramento Valley, a Common Ground-Dove was discovered at Elk Creek, Glenn 4 Jul (ph. Sarah MacLean). Two Yellowbilled Cuckoos included Del Norte's first along the Smith River 3-4 Jun (TKz, Casey Ryan, ph. LuB, ADB) and the third for Arcata Marsh, Humboldt 27-31 Jul (AxL, m.ob.). Short-eared Owls have been found in Mono during the nesting season on a few occasions over the past several decades, suggesting probable breeding. However, 2 fledglings at Mono Lake C.P. 19 Jul (LHg) provided the first confirmed breeding record since 1933.

A total of eight Black Swift nests at Burney Falls, *Shasta* 20 Jul (EE) was not unusual for this traditional nesting location. What was unusual, however, was that seven nests were visible in one scope view! Each nest contained a large nestling, one of which was being brooded by an adult. White-throated Swifts are scarce in *Sierra*, so one well-described bird over Sierra City 28 Jun (†RC) was noteworthy. *Del Norte's* first Broad-billed Hummingbird (p.a.) visited a Darlingtonia bog at Smith River R.A. 27 Jun (†Casey Ryan, ADB). As most of our records of vagrant hummingbirds come from feeders, this record is especially notable.

Lewis's Woodpeckers are scarce anywhere along the immediate coast, where they sometimes occur fall-spring. Therefore, one in Tennessee Valley, *Marin* 4 Jun (Jeff Kimball, Mark Forney) was completely unexpected. Following last winter's invasion, Acorn Woodpeckers persisted in S.F. through the period, with at least 4 in Lafayette Park plus 10 or more in Golden Gate Park, where nesting was confirmed 27 Jun (Dylan Hayes), providing the first breeding record for *S.F.* A Nuttall's Woodpecker, evidently a post-breeding wanderer, at Doyle, *Lassen* 23 Jul (Denise Riddle), was in the e. part of the county, well outside areas of normal occurrence for this non-migratory species.

Our only Least Flycatcher was along the Smith River, *Del Norte* 2-7 Jun (LuB, m.ob.). Likewise, our only Eastern Phoebe was in *Del Norte*, at the Crescent City Airport 1 Jun (LuB). Eastern Kingbirds were more plentiful, with 8 birds reported, including *Stanislaus*'s first at Faith Ranch 14 Jul (ph. Gary Zahm) and two pairs at the traditional breeding site at Blue Lake, *Lassen* 22 Jul (ph. EE). Other singles were at P.R.N.S., *Marin* 8 Jun (Mark Forney), Point Saint George, *Del Norte* 9 Jun (ADB), and Pescadero Marsh, *San Mateo* 25 Jun (GrH).

A White-eyed Vireo at Campbell Cove 7-8 Jun (ph. DN, v.r. ESH, m.ob.) furnished the first record for Sonoma. Continuing from spring, the Bell's Vireo w. of Corcoran, Kings lingered until 7 Jun (MSy); it was the only one reported. Our only Yellow-throated Vireo, a singing male, was found along Virginia Creek, Mono 17 Jun (KNN). Plumbeous Vireos included a remarkable 12 (six pairs on territory) in a small stringer of aspens in Wildrose Canyon, Mono 12 Jun (CMc) and a single bird outside of breeding range at Klamath Glen, Del Norte 14 Jun-29 Jul (LuB, KBu). Our only Philadelphia Vireo visited P.R.N.S., Marin 7 Jun (Everett Clark). Five Redeyed Vireos included singles at Redwood N.P., Humboldt 5 Jun-30 Jul (GSL, LPL, ph., KBu, COg), P.R.N.S., Marin 7 Jun (Alex Merritt), Cache Creek 8 Jun (Yolo's fifth; Brent Campos), Klamath Glen, Del Norte 9-13 Jun (LuB, JCS), and Eureka, Humboldt 10 Jun (TWL).

About 10 ad. Purple Martins associating with two occupied nest cavities at the Bonny Doon Ecological Reserve throughout the period (AMR, GJS, Craig Fosdick et al.) established the first nesting record for *Santa Cruz* since 2001. Similarly, a string of sightings in Jul, including that of 2 ads. feeding 2 juvs. at Pillar Point Harbor 8 Jul (RSTh), indicated the first successful nesting in the *San Mateo* portion of the Santa Cruz Mountains in 20 years. Four Swainson's Thrushes singing along a 3-km stretch of Mono Creek ne. of Edison Lake, *Fresno* 3 Jul (JTz) suggested probable breeding. Although this species was thought to have bred widely along the w. slope of the Sierra Nevada historically, there are no specific breeding records for that county.

THRASHERS THROUGH WARBLERS

A Gray Catbird was mist-netted at Bolinas, Marin 16 Jun (Renee Cormier), and another was found singing 22 Jun just 16 km n. of Bolinas at White House Pool, Marin (DvH), raising the possibility that one bird was responsible for both observations. An out-of-range Sage Thrasher was at the Crescent City Airport, Del Norte 9 Jun (Sky Lloyd, ph. LuB). Our only Brown Thrasher was photographed at P.R.N.S. 7 Jun (Mark Butler, ph. Everett Clark). Three of our 4 Ovenbirds were on F.I. 6-21 Jun (Point Blue); the fourth was at P.R.N.S. 13 Jun (Alex Merritt, ph. Everett Clark, W. Durrel Kapan). Five Black-and-white Warblers, all in coastal or S.F. Bay counties, furnished a typical seasonal total. A Prothonotary Warbler at P.R.N.S. 11-19 Jun (DFr, ph. Mark Forney) was a good find. A Tennessee Warbler was banded on F.I. 7 Jun (Point Blue), and another was found singing at Campbell Cove, Sonoma 9 Jun (DN, DFr). Hooded Warblers were at P.R.N.S. 19 Jun (Mark Sawyer) and at H.R.S. 25 Jun (RJR). Eight American Redstarts included 2 inland birds, one at Oro Loma, Fresno 9 Jun (JSy) and another at Devils Postpile N.M. 19 Jul (ph. Bruce Aird; Madera's third). Eight of 12 Northern Parulas were in Humboldt. A singing male parula provided Lake's second record at Corinthian Bay 16 Jun (JRW), and another singer supplied a notable mid-summer record for Yolo, at the Sacramento Bypass 29 Jun (SCH). Five Magnolia Warblers produced a total twice our summer average. A Blackburnian Warbler on F.I. 5 Jun (Point Blue) was our first in summer since 2009. Summer reports of Yellow Warblers on the C.V. floor seem to be increasing, so birds in good riparian habitat at the Sacramento River N.W.R., Pine Creek Unit, Butte 6 Jun (FrO et al.) and Capay Unit, Glenn 15 Jun (MMu) provided encouragement that this species may someday re-colonize this part if its historical range. Seven Chestnut-sided Warblers provided a good showing, as we average fewer than 5 per season. Blackpoll Warblers were at Ardenwood Farm, Alameda 23-24 Jun (Marty Morrow, ES, ph. m.ob.) and on F.I. 4-7 Jul (Point Blue), and our only Palm Warbler was at Campbell Cove, Sonoma 9 Jun (DN, DFr).

SPARROWS THROUGH BLACKBIRDS

Wintering Savannah Sparrows often linger into May and begin singing before they leave grassland habitats in the C.V. and other areas in our Region. Records of singing birds (away from the delta/coastal and high Sierra habitats) in late Jun are unusual. Therefore, observations of Savannah Sparrows singing in grassland habitats along the Pajaro River, Santa Cruz 17 Jun (NLv) and at the Sacramento N.W.R., Glenn 27 Jun (Sarah MacLean) are intriguing. A White-throated Sparrow at Petaluma, Sonoma 4 Jul (Sherri Oster) provided a rare late summer record. Other lingering Zonotrichia sparrows included a White-crowned Sparrow along the Pajaro River near the Santa Clara county line, Santa Cruz 17 Jun (NLv) and a Golden-crowned Sparrow on San Bruno Mountain, San Mateo 23 Jun–2 Jul (Doug Allshouse). Summer Tanagers were at Acampo, San Joaquin 5 Jun (DGY et al.), at Ed Levin C.P., Santa Clara 22 Jun (WGB), and at Coyote Creek Field Station, Santa Clara 29 Jul (ph. Josh Scullen). A female Scarlet Tanager at Campbell Cove 1 Jun (ph. DN, PBC) provided Sonoma its first record. A stunning ad. male Scarlet Tanager was at Ardenwood Farm 19-21 Jun (Chingling Tien, ph. Bill Chen, ph. Bob Toleno, m.ob.), only the second of this species for Alameda. Fifteen Rose-breasted Grosbeaks and 5 Indigo Buntings produced totals for both species well below our averages (26 and 11, respectively). A Blue Grosbeak singing along Pescadero Creek, Santa Cruz 3 Jul (AMR) was very close to a known breeding location in Santa Clara, suggesting that this site may yet produce a first confirmed breeding record for Santa Cruz. The Tricolored Blackbirds found in spring along the Carson River, Alpine were still present 3 Jun (Nancy Overholtz); unfortunately, we have no later reports and no confirmation that these birds bred.

Cited observers (county subregional editors in boldface): Ken Able (Lassen), David P. Arsenault, Alan D. Barron, Dave A. Bengston, Murray Berner (Napa), William G. Bousman (Santa Clara), Lucas Brug, Ken Burton, Rita Carratello, Peter B. Colasanti, Daryl Coldren (Humboldt and Del Norte), Chris Conard (Sacramento), Rudy C. Darling (Nevada), Jeff N. Davis (Madera), Malia DeFelice, Bruce E. Deuel (n. C.V. counties), Colin P. Dillingham (Plumas and Sierra), Cedric Duhalde, Leo J. Edson, Alan M. Eisner, Elias Elias, Rob Fowler (RbF; Humboldt and Del Norte), Dea Freid, Steve Gerow (Santa Cruz), David Hamilton, Steve C. Hampton, John H. Harris (Stanislaus), Garth Harwood, Floyd Hayes, Cliff Hawley, Debbie House, Lisa Hug, Eugene S. Hunn, John E. Hunter (Trinity), Richard G. Jeffers, Logan Kahle (Contra Costa and San Francisco), Robert J. Keiffer (Mendocino), Tony Kurz, Alexandra Lamb, Kimya L. Lambert, Sarah Lane, Robin L. C. Leong (Solano), Tom W. Leskiw, Gary S. Lester, Lauren P. Lester, Nick Levendosky, John Lewis, Jim Lomax, Michael J. Mammoser, Dan J. Maxwell (Marin), Chris McCreedy, Tristan McKee, Peter J. Metropulos (San Mateo), Dominik Mosur (San Francisco), Mary Muchowski, Dan Nelson, Kristie N. Nelson (Mono), Chet Ogan, Frances Oliver (San Joaquin), Ed Pandolfino (Placer), Ryan Phillips, Linda Pittman, Gary W. Potter (Fresno), Robert C. Power, Harold M. Reeve, Robert J. Richmond (Alameda), Alex M. Rinkert, Don Roberson (Monterey), Michael M. Rogers (MMR), Michael Rogner (MRg), Diane Rose, Steve Rose, Stephen C. Rottenborn, James Rowoth, Ruth A. Rudesill (Sonoma), Jeff Seay (Kings), Daniel Singer, Mark Stacy, John C. Sterling (Modoc, Alpine, Calaveras, and Yolo), Tim Steurer (El Dorado and Amador), Gary J. Strachan, Emilie Strauss, Steve J. Stump, David Styer, Brian L. Sullivan, Steven D. Summers (Tulare), Ronald S. Thorn, Jim Tietz (EL), Steve Umland (Tuolumne), Kent Van Vuren (Merced), Jerry R. White (Lake), Gary Woods, David G. Yee, Bob & Carol Yutzy (Shasta). Many more observers are not specifically cited; all are appreciated.

Michael M. Rogers (WATERFOWL TO QUAIL, HERONS TO SHORE-BIRDS), 499 Novato Avenue, Sunnyvale, California 94086 (m.m.rogers@comcast.net)

Jeff N. Davis (DOVES TO THRUSHES), 11238 N. Via Trevisio Way, Fresno, California 93730 (jdavis@colibri-ecology.com) Ed Pandolfino (THRASHERS TO FINCHES), 1328 49th Street, Sacramento, California 95819 (erpfromca@aol.com) Stephen C. Rottenborn (LOONS TO FRIGATEBIRDS, LARIDS TO ALCIDS), 983 University Avenue, Building D, Los Gatos, California 95032 (srottenborn@harveyecology.com)



Guy McCaskie Kimball L. Garrett

SPRING

ridge of high pressure ruled southern California in March, with temperatures well above normal and precipitation well below normal away from the southeastern deserts. Temperatures were only slightly above normal in April, but rainfall was still well below average. By May, temperatures cooled to below average, and although there was above-average rainfall in much of the Region that month, we still headed into summer having experienced a fourth consecutive exceptionally dry rainy season. Offshore, sea surface temperatures were above normal and continuing to warm. Avian highlights were few this season, with a generally lackluster migration and spring vagrant season. Most noteworthy was a Nazca Booby at Anacapa Island in May (only the third to be found north of Mexico), and three Common Black Hawks on the southern deserts in March and April. Lesser but still significant rarities included a Cape May Warbler, a Cassin's Sparrow, and an

unprecedented early April Mississippi Kite.

Abbreviations (both reports): C.L. (China Lake Naval Air Weapons Station, extreme ne. Kern); E.A.F.B. (Edwards A.F.B., se. Kern); F.C.R. (Furnace Creek Ranch, Death Valley N.P., Inyo); G.H.P. (Galileo Hill Park in extreme e. Kern); N.E.S.S. (n. end of the Salton Sea, Riverside); P.M.N.A.S. (Point Mugu Naval Air Station, Ventura); P.P. (Piute Ponds on Edwards A.F.B., ne. Los Angeles); S.E.S.S. (s. end of the Salton Sea, Imperial); S.F.K.R.P. (South Fork Kern River Preserve near Weldon, Kern); S.J.W.A. (San Jacinto Wildlife Area near Lakeview, Riverside); V.C.G.P. (Ventura County Game Preserve near Point Mugu Naval Air Station, Ventura). Because most rarities in s. California are seen by multiple observers, only the observer(s) initially finding and/or identifying the bird are included. Documentation for species on the California Bird Records Committee review list (see www.californiabirds.org) is forwarded to the committee and archived at the Western Foundation of Vertebrate Zoology in Camarillo.

WATERFOWL THROUGH HERONS

Each spring, migrant waterbirds wintering in the Gulf of California follow a non-stop route that takes them northwestward across the Anza Borrego Desert and the lower parts of the Laguna Mountains to the Pacific, probably across Los Angeles, as suggested by 40 Brant flying westward over El Monte 12 Apr (LSc), stopping only when forced to the ground by storms. Flocks of 45 Brant on Fig Lagoon near Seeley, Imperial 11 Apr (GMcC) and 15 on Lake Henshaw, San Diego 21 Mar (RA), along with up to 10 together at multiple locations at the Salton Sink, Imperial/Riverside, and the general area around Borrego Springs, San Diego between 28 Mar and 20 May, were all such storm-grounded birds. However, 4 at Bishop, Inyo 21 May (JJB) had erroneously fol-

Southern California

lowed a more easterly route that placed them on the e. side of the Sierra Nevada. The latest of the wintering Eurasian Wigeons were single birds at Kern N.W.R., Kern through 18 Apr (AMS), the S.J.W.A. through 26 Apr (CMcG), and at Hansen Dam, Los Angeles through 29 Apr (JR). Long-staying Harlequin Ducks at Estero Bluffs S.P., San Luis Obispo through 30 May (TME) and at Avila Beach, San Luis Obispo through 16 May (JCop) were expected to remain through the summer. Single Surf Scoters at Borrego Springs 8 Apr (PA) and on the Salton Sea at Salt Creek 23 Apr (RMcK) and a White-winged Scoter at Lancaster, Los Angeles 24 Apr-4 May (CH) were inland, where rare, but much rarer was a Black Scoter at Borrego Springs 8 Mar (PA). In addition, the total of 15 Black Scoters along the coast of San Luis Obispo was far more than expected, with the latest being one near Cayucos 25-30 May (TME). A Long-tailed Duck was inland at Desert Center, Riverside 13-14 Mar (BS), and another at Kern N.W.R. 14-25 Mar (BO'C) was only the fourth to be found in Kern; on the coast, single birds were at Morro Bay, San Luis Obispo 21 Mar-20 May (JM) and Pismo Beach, San Luis Obispo 2-9 May (P&JC).

A Sooty Grouse at Silver Canyon 10 Apr (SLS) was in the White Mountains, Inyo, where few have been reported in recent years. Single Pacific Loons on Hesperia Lake, San Bernardino 2-22 May (MWo), at N.E.S.S. 10 (CMcG) & 29 Mar (CMcG), at Borrego Springs 23 Apr-1 May (WTH), and at Fig Lagoon 11 Apr (GMcC) were all well inland, where rare. A Red-necked Grebe on the coast at P.M.N.A.S. 20 Mar (DPe) was the only one reported. A Laysan Albatross at the 9-Mile Bank off San Diego 3 & 16 May (DPo) was only the third live bird ever recorded in San Diego waters. A Laysan Albatross was captured well inland at a construction site in Palm Desert, Riverside 20 Mar and taken to San Pedro for treatment and release. Noteworthy procellariids reported by PEL from cruise-